

Development of ^{166}Ho -zoledronate as a bone marrow ablative agent

Mirsaeed Nikzad¹, Amir Reza Jalilian^{2*}, Simindokht Shirvani-Arani², Ali Bahrami Samani², Hamid Golchobian¹

¹Department of Inorganic Chemistry, University of Mazandaran, Babol-sar, Iran

²Nuclear Science and Technology Research Institute (NSTRI), Tehran, Iran

Received: Feb 14, 2016, Revised: Feb 27, 2016, Accepted: Mar 2, 2016

Abstract

In this study, production, quality control and biodistribution studies of ^{166}Ho -zoledronate have been presented as a possible bone marrow ablative agent. Ho-166 chloride was produced by thermal neutron irradiation of natural $^{165}\text{Ho}(\text{NO}_3)_3$ samples. ^{166}Ho -zoledronate complex was prepared by adding the desired amount of zoledronate solution (0.2 mL, 150 mg/ml in 1 M NaOH) to appropriate amount of the $^{166}\text{HoCl}_3$ solution. Radiochemical purity of the complex was monitored by instant thin layer chromatography (ITLC). Stability studies of the complex in the final preparation and in the presence of human serum were performed up to 48 h. The biodistribution of ^{166}Ho -zoledronate and $^{166}\text{HoCl}_3$ in wild-type mice was checked up to 72 h. ^{166}Ho -zoledronate complex was prepared in high radiochemical purity (> 99%, ITLC) and specific activity of 4.4 GBq/mmol. The major accumulation of radiolabelled complex was observed in the bone tissue. These findings suggest ^{166}Ho -zoledronate has can be a possible candidate for bone marrow ablation in patients with multiple myeloma.

Keywords: Bone marrow ablation, Ho-166, Zoledronate, biodistribution

Pharm Biomed Res 2016; 2(1): 14-22

DOI: 10.18869/acadpub.pbr.2.1.14

Introduction

Over the past 30 years, bisphosphonates were established as an effective new drug class for bone pain palliation in patients with different complicated malignancies such as prostate, breast and lung cancer (1). Bisphosphonates have strong affinity for calcium phosphates and for hydroxyapatite and attach preferentially to bone with increased avidity for areas of osteoblastic and osteoclastic activity (2,3).

Nowadays, researches have shown that skeletal targeted radiotherapy in patients

with multiple myeloma can improve the response rate in phase I and II trials, with promising long-term survival data (4). Several bone-seeking radiopharmaceuticals have been developed for bone marrow ablation in multiple myeloma and other haematological malignancies (5).

Samarium-153-ethylene diamine tetramethylene phosphonate (^{153}Sm -EDTMP) is now considered as an efficient radiopharmaceutical for bone pain palliation (6). Various other radiopharmaceuticals such as rhenium-

* E-mail: E-mail:ajalili@aeoi.org.ir

¹⁸⁶Re-hydroxyethylidenediphosphonic acid (¹⁸⁶Re-HEDP) (7), ¹⁶⁶Ho-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetramethylene-phosphonate (¹⁶⁶Ho-DOTMP) (8) and Tin-117m (9) have also been investigated for targeted radiotherapy of bone malignancies.

In order to give sufficient dose to the bone marrow, radionuclides with higher beta particle energy are recommended for bone marrow ablation. ¹⁶⁶Ho with a half-life of 26.8 h decays with two principal β^- emissions (1.85 MeV (51%) and 1.77 MeV (48%)), and three photon energies (80.58 MeV (6.56%), 1379.44 (0.92%) and 1581.83(0.18%)) (10) is an excellent radionuclide for bone marrow ablation due to the high energy of the β^- particles. The short physical half-life permits delivery of high-dose irradiation to peripheral blood stem cells during chemotherapy and reinfusion of cryopreserved within 6-10 d (8).

Various therapeutic possible ¹⁶⁶Ho bone-seeking agents have been reported such as ¹⁶⁶Ho-DOTMP (8), ¹⁶⁶Ho-EDTMP (5) and ¹⁶⁶Ho-APDDMP (11). The presence of extra amounts of bisphosphonate ligand in the final formulation is inevitable in all mentioned agents which can impose toxicity concerns especially for non-pharmaceutical ligands, however the use of a pharmaceutical ligand such as zoledronate with known toxic dose and biological behaviour and availability can be a superiority. However, the search for the development of new ¹⁶⁶Ho-bisphosphonate ligands with higher stability, better pharmacokinetics and lower unwanted tissue uptakes (liver and gastrointestinal tract) is still ongoing.

there are two separate categories for beta emitters used in clinics, i.e. bone pain palliative and bone ablative agents. The difference is mostly based on the range

of beta particle traveling in the tissues. For instance Sm-153 has a lower range and usually does not reach the bone marrow tissue and is just reducing the bone proliferative cancerous cells, while Ho-166 with longer range of beta emission is really affecting the bone marrow and its radiopharmaceuticals are intended for irradiation of bone marrow tissue in bone marrow cancers.

A number of bisphosphonates such as clodronate, etidronate, pamidronate, zoledronate and ibandronate are commercially available in some countries. Among the mentioned bisphosphonates, zoledronate (Fig. 1) is a potent agent which inhibits osteoclast-mediated bone-resorption (12). In inhibiting the bone-resorption, zoledronate is approximately 1,000 times stronger than etidronate (13). Moreover, the binding affinity of zoledronate for bone is 10 orders of magnitude higher than pamidronate (14).




Figure 1 Chemical structure of zoledronic acid

Treatment of post-menopausal women with zoledronate has demonstrated normalization of the rate of bone turnover, significant increase bone mineral density of the spine, hip, wrist and total body, and significant reductions in the risk of vertebral

(spine) fractures, wrist fractures, hip fractures, and all non-vertebral fractures (15). In this work, ¹⁶⁶Ho-zoledronate was prepared as a new complex for bone marrow ablation and biodistribution studies of this complex was investigated in wild-type mice up to 72 h.

Materials and methods

Materials and instruments

Natural holmium nitrate with purity of >99.99% was obtained from ISOTECH Inc (USA). Production of ¹⁶⁶Ho was performed using ¹⁶⁵Ho(n,γ)¹⁶⁶Ho nuclear reaction in Tehran Research Reactor (5 MW). The precursor zoledronate hydrochloride with Pharmaceutical grade was a gift from Sobhan Co., Tehran Iran, with > 99% purity. Radio-chromatography was performed by counting of ITLC-SG and/or Whatman papers using a thin layer chromatography scanner, Bioscan AR2000 (Paris, France). ITLC-SG was purchased from UK. Other chemicals were purchased from Sigma-Aldrich. A high purity germanium (HPGe) detector coupled with a Canberra™ (model GC1020-7500SL) multichannel analyzer and a dose calibrator ISOMED 1010 (Dresden, Germany) were used for counting distributed activity in mice organs. Calculations were performed based on the 80.6 keV peak for ¹⁶⁶Ho. Animal studies were performed in accordance with the United Kingdom Biological Council's Guidelines on the Use of Living Animals in Scientific Investigations, 2nd edn. The wild-type mice weighing between 20-25 g were acclimatized at proper rodent diet .

Production and quality control of ¹⁶⁶HoCl₃ solution

The desired amount (about 100 μg) of natural ¹⁶⁵Ho(NO₃)₃ was irradiated at a thermal neutron flux of 4×10¹³ n.cm⁻².s⁻¹. The irradiated target was dissolved in 200 μl of 1.0 M HCl, to prepare ¹⁶⁶HoCl₃. The solution was filtered through a 0.22 μm filter (Millipore, Millex GV, USA). The radionuclidic purity of the solution was checked using beta liquid scintillation as well as HPGe spectroscopy. Also the radiochemical purity of the ¹⁶⁶HoCl₃ was studied using instant thin layer chromatography (ITLC) method.

Radiolabeling of zoledronate with ¹⁶⁶HoCl₃

A stock solution (50 mL) of zoledronate was prepared by dissolving in 1 M NaOH and diluted to the appropriate volume with ultra-pure water, in order to produce a solution of 20 mg/ml. For labelling, an appropriate amount of the ¹⁶⁶HoCl₃ solution, containing the required activity was added to the desired amount of zoledronate solution (0.2 mL). The complex was maintained at 40°C for 2 hours. The final solution was passed through a 0.22 μm membrane filter and the pH was adjusted to 7-8.5 with 0.05 M phosphate buffer (pH 5.5). The radiochemical purity was determined using Whatman 3 MM chromatography paper or ITLC-SG, eluted with NH₄OH (56%): methanol (100%): water (100%) (0.2:2:4; v/v/v) mixture.

Stability studies

The stability of the complexes stored at room temperature (25 °C ambient) and in human serum at 37 °C at different time intervals were checked by determining the radiochemical purity of the complex by paper chromatography

in $\text{NH}_4\text{OH}:\text{MeOH}:\text{H}_2\text{O}$ (2:20:40) system.

In vitro stability of ^{166}Ho -zoledronate in presence of human serum

Final solution (200 μCi , 50 μL) was incubated in the presence of freshly prepared human serum (300 μL) (Purchased from Iranian Blood Transfusion Organization, Tehran, Iran) and kept at 37 °C for 2 h. Trichloroacetic acid (10%, 100 μL) was added to a portion of the mixture (50 μL) every 30 min and the mixture was centrifuged at 3000 rpm for 5 min. The debris was decanted from the supernatant. The stability was determined by performing frequent ITLC analysis of supernatant using above mentioned ITLC system.

Hydroxyapatite binding assay

The hydroxyapatite binding assay was performed according to the procedure described previously (16), with only a slight modification. In brief, to vials containing 1.0, 2.0, 5.0, 10.0, 20.0 and 50.0 mg of solid hydroxyapatite, 2mL of saline solution of pH 7.4 were added and the mixtures were shaken for 1h. Then, 50 μL of the radioactive preparation (containing 100-300 μCi) was added and the mixtures were shaken for 24 h at room temperature. The suspensions were centrifuged, and two aliquots of the supernatant liquid were taken from each vial and the radioactivity was measured with a well-type counter. Test experiments were performed using a similar procedure, but in the absence of hydroxyapatite. The percentage binding of ^{166}Ho -labeled compound to hydroxyapatite (HA) was calculated according to $\text{HB} = 1 - \text{A}/\text{B} \times 100$, where A is the mean radioactivity value of the supernatant sample under study

and B is the mean total value of whole activity used.

Biodistribution of ^{166}Ho cation and ^{166}Ho -zoledronate in wild-type mice
50-100 μL of $^{166}\text{HoCl}_3$ and final ^{166}Ho -zoledronate solution with 200 μCi radioactivities was injected intravenously to mice through their tail vein. The animals were sacrificed at the exact time intervals (2, 4, 24, 48, 72h) post injection the percent injected dose per gram (%ID/g) of different organs was calculated based on the measurements of HPGe detector. Three animals were used for each time interval.

Results

Radionuclide production

The radionuclide was prepared in a research reactor with a range of specific activity between 3 to 5 GBq/mg. After counting the samples on an HPGe detector for 5 min, two major photons (5.4% of 80.68 keV and 0.9% of 1379.94 keV) were observed. Radionuclide purity was higher than 99.96%. The radiochemical purity of the $^{166}\text{HoCl}_3$ was checked using two solvent systems for ITLC (A: 10 mM DTPA pH 4 and B: ammonium acetate 10% : methanol (1 : 1)).

Radiolabeling of zoledronate with $^{166}\text{HoCl}_3$

In order to obtain maximum complexation yield, several experiments were carried out by varying different reaction parameters such as ligand concentration, pH, and reaction time. The radiochemical purity of more than 99% was achieved at pH 7-8 and 40-50 °C with 40:1 ligand molar ratio to calculated holmium content. The reaction mixture was incubated at room

temperature for different time periods and 120 min incubation was found to be adequate to yield maximum complexation. The best ITLC mobile phase was considered Whatman 3 MM paper using NH₄OH: MeOH: H₂O (0.2:2:4). In the chromatography system of choice, free holmium cation elutes to the R_f of 0.05 while the radiolabelled bis-phosphonate complex migrates to higher R_fs, i.e. 0.9 (Fig. 2)

Stability studies

The stability of the complex prepared under optimized reaction conditions was studied at room and in presence of human serum at 37 °C. It was observed that the complex was > 95% stable both at room temperature and in presence of human serum at 37 °C even after 48 h.

Hydroxy apatite assay

The assay demonstrated high capacity binding for ¹⁶⁶Ho-Zoledronate to hydroxy apatite. Even at 2-3 mg amount of HA, more than 85% binding was observed, while at 5 mg HA used > 92% of binding was obtained. Compared to the other radiolanthanide zoledronate complex, i.e., ¹⁵³Sm-zoledronate, the holmium complex demonstrates higher hydroxy apatite binding, while Sm complex reaches the 90% affinity at 15 mg of hydroxyapatite (16), ¹⁶⁶Ho-zoledronate, retains this binding at 2-3 mg which is far better *in vitro* data for a bone seeking bis-phosphonate (Fig. 3).




Figure 2 ITLC chromatograms of ¹⁶⁶HoCl₃ (left) and ¹⁶⁶Ho-Zoledronate solution (right) using Whatman 3 MM eluted with NH₄OH: MeOH: H₂O (0.2:2:4)




Figure 3 ^{166}Ho -Zoledronate binding assay to hydroxyapatite (1-20 mg)

Biodistribution of ^{166}Ho cation and ^{166}Ho -zoledronate in wild-type rats

The animals were sacrificed by CO_2 asphyxiation at selected time points after tracer injection. Dissection began by drawing blood from the aorta followed by removing the heart, spleen, muscle, bone, kidneys, liver, intestine, stomach and lungs samples. The tissue uptakes were calculated as the percent of area under the curve of the related photo peak per gram of tissue (%ID/g). For $^{166}\text{Ho}^{3+}$ cation, the radioactivity was mainly located in the liver, kidney and bone. The free cation is soluble in water and it can be excreted via the urinary tract (Fig. 4).

Discussion

Radiolabelling of the complex using the carrier added Ho-166 produced in the research reactor was consistent with other related radiolanthanides as reported previously (4-7). Variations in complexation yields with respect to the zoledronate:holmium molar ratio as shown in Table 1 led to optimization of ligand concentration used. At pH 7-8 and 40 °C the best saturated molar ratio was observed for 50:1 ratio, while a major drop in radiochemical purity is observed in reducing 20 to 10. The stability of the complex at least in vitro was encouraging to initiate the animal studies according to the ethical animal codes.




Figure 4 Percentage of injected dose per gram of $^{166}\text{HoCl}_3$ in wild-type rat tissue after 2, 4, and 24 h post injection

Table 1 Complexation yields of ¹⁶⁶Ho- Zoledronate at different [ligand]:[metal] ratios

| [ligand]:[metal] ratios | Radiochemical purity (%) |
|-------------------------|--------------------------|
| 1:1 | 4 |
| 1:10 | 24 |
| 1:20 | 43 |
| 1:30 | 67 |
| 1:50 | 99 |
| 1:75 | 99 |

On the other hand, hydroxy apatite binding was also higher than 95% which is in accordance with in vitro data obtained for other bisphosphonate complexes (4-7).

Approximately 1.5% of the ¹⁶⁶Ho cation accumulates in the liver after 2 h which is comparable with many other radio-lanthanides mimicking calcium cation accumulation. Ho-166 activity is washed out as a water soluble free cation through kidney in 24 h. Ho-166 cation is present in other tissues are negligible. The distribution of injected dose in wild-type rat organs up to 72 h after injection of ¹⁶⁶Ho-zoledronate was also determined. As it can be seen in Figure 5, the major portion of the injected activity of ¹⁶⁶Ho-zoledronate was accumulated in bones as expected for bone-avid radiopharmaceuticals. For better understanding the ratios of target:non target for this complex in various time intervals Figure 6 demonstrates the increasing the uptake




Figure 5 Percentage of injected dose per gram of ¹⁶⁶Ho-zoledronate in wild-type rat tissue after 2, 4, 24, and 48h post injection




Figure 6 Comparative Bone:Blood, Bone:liver, Bone:Kidney activity ratios for ^{166}Ho -zoledronate in 2-72 h

ratio with increasing time up to 72 h. Compared to other reported Ho-166 labeled bisphosphonates such as ^{166}Ho -DOTMP (18), the bio distribution of this complex looks superior, however the slightly higher liver uptake in this case is a drawback. On the other hand non-cyclic bisphosphonate complexes such as ^{166}Ho -pamidronate have shown far worse biodistribution data leading to higher kidney and liver uptake possibly due to in vivo stability (19). The only non-cyclic bisphosphonate showing good performance in animal data has been ^{166}Ho -EDTMP (20) with almost acceptable stability and bone targeting. The aim of radionuclide therapy is complete ablation of the target with minimal radiation exposure to other organs. Target to non-target ratio is the most important parameter in the selection of radiopharmaceuticals for therapy of choice for each malignancy. Since ^{166}Ho -zoledronate was produced as a bone marrow ablative agent, red marrow to other tissue dose must be determined.

Conclusion

^{166}Ho -zoledronate complex was prepared in high radiochemical purity (>99 %, ITLC) and specific activity of 4.4 GBq/mmol. The complex demonstrated significant stability at room temperature and in presence of human serum at 37 °C even after 48 h. The final preparation was administered to wild-type rats and biodistribution of the complex was checked 2-72 h post injection showing major accumulation of the complex in the bone tissue. The highest absorbed dose for ^{166}Ho -zoledronate is observed in bone tissue. Compared to the other radiolanthanide zoledronate complexes, ^{166}Ho -zoledronate demonstrates higher hydroxy apatite binding and also better bone:liver ratio. According to the results, ^{166}Ho -zoledronate has considerable characteristics and can be a good candidate for bone marrow ablation in patients with multiple myeloma.

References

1. Palma E, Correia JDG, Campello MPC, Santos I, Bisphosphonates as radionuclide carriers for imaging or systemic therapy. *Mol BioSyst* 2011;7:2950-66.
2. Fleisch H, Russell RG, Francis MD, Diphosphonates inhibit hydroxyapatite dissolution in vitro and bone. *Science* 1969; 165:1262-64.
3. Majkowska A, Neves M, Antunes I, Bilewicz A, Complexes of low energy beta emitters ⁴⁷Sc and ¹⁷⁷Lu with zoledronic acid for bone pain therapy. *Appl Radiat Isot* 2009;67:11-3.
4. Giralt S, Bensinger W, Goodman M, Podoloff D, Eary J, Wendt R, et al. ¹⁶⁶Ho-DOTMP plus melphalan followed by peripheral blood stem cell transplantation in patients with multiple myeloma: results of two phase I/II trials. *Blood* 2003;102:2684-91.
5. Pedraza-López M, Ferro-Flores G, de Murphy CA, Tendilla JI, Villanueva-Sánchez O. Preparation of (166)Dy/(166)Ho-EDTMP: a potential in vivo generator system for bone marrow ablation. *Nucl Med Commun* 2004; 25:615-21.
6. Abruzzese E, Iuliano F, Trawinska MM, Di Maio M. ¹⁵³Sm: its use in multiple myeloma and report of a clinical experience. *Expert Opin Investig Drugs* 2008;17:1379-87.
7. Wiesmann A, Einsele H, Kanz L, Dohmen BM. Complete remission after myeloablation with bone-seeking ¹⁸⁶Re-HEDP and high-dose melphalan followed by autologous stem cell transplantation in a patient with chemorefractory multiple myeloma. *Bone Marrow Transplant* 2005; 36:89-90.
8. Breitz HB, Wendt III RE, Stabin MS, Shen S, Erwin WD, Rajendran JG, et al. ¹⁶⁶Ho-DOTMP radiation-absorbed dose estimation for skeletal targeted radiotherapy. *J Nucl Med* 2006;47:534-42.
9. Srivastava SC, Atkins HL, Krishnamurthy GT, Zanzi I, Silberstein EB, Meinken G, et al. Treatment of metastatic bone pain with tin-117m Stannic diethylenetriaminepentaacetic acid: a phase I/II clinical study. *Clin Cancer Res* 1998; 4:61-8.
10. Calhoun JM, Cessna JT, Coursey BM, Hoppes DD, Schima FJ, Unterweger MP. Standardization of holmium-166 by the CIEMAT/NIST liquid scintillation efficiency-tracing method. *Radioact Radiochem* 1992; 2:38-45.
11. Zeevaart JR, Jarvis NV, Louw WK, Jackson GE. Metal-ion speciation in blood plasma incorporating the tetrakisphosphate, N,N-dimethylenephosphonate-1-hydroxy-4-aminopropylidenediphosphonate (APDDMP), in therapeutic radiopharmaceuticals. *J Inorg Biochem* 2001;83:57-65.
12. Astrand J, Aspenberg P. Topical, single dose bisphosphonate treatment reduced bone resorption in a rat model for prosthetic loosening. *J Orthop Res* 2004;22:244-9.
13. Alendronic Acid information from Drugs Update, Available at "http://www.drugsupdate.com/generic/view/89/Alendronic-Acid".
14. Kanis JA, Gertz BJ, Singer F, Ortolani S. Rationale for the use of alendronate in osteoporosis. *Osteoporos Int* 1995;5:1-13.
15. Black DM, Cummings SR, Karpf DB, Cauley JA, Thompson DE, Nevitt MC, et al. Randomised trial of effect of alendronate on risk of fracture in women with existing vertebral fractures. Fracture Intervention Trial Research Group. *Lancet* 1996;348: 1535-41.
16. Neves M, Gano L, Pereira N, Costa MC, Costa MR, Chandia M, Rosado M, Fausto R. Synthesis, characterization and biodistribution of bisphosphonates Sm-153 complexes: correlation with molecular modeling interaction studies. *Nucl Med Biol* 2002; 29:329-38.
17. Talwer, GP, Srivastava, LM. Textbook of biochemistry and human biology.
18. Bagheri R, Jalilian AR, Bahrami-Samani A, Mazidi A, Ghannadi-Maragheh M. Production of Holmium-166 DOTMP: A promising agent for bone marrow ablation in hematologic malignancies. *Iran J Nucl Med* 2011;19:12-20.
19. Fakhari A, Jalilian AR, Yousefnia H, Johari-Daha F, Mazidi M, Khalaj A. Development of ¹⁶⁶Ho-pamidronate for bone pain palliation therapy. *J Radioanal Nucl Chem* 2015;303:743-50.
20. Bahrami-Samani A, Bagheri R, Jalilian AR, Shirvani-Arani S, Ghannadi-Maragheh M, Shamsaee M. Production, quality control and pharmacokinetic studies of ¹⁶⁶Ho-EDTMP for therapeutic applications. *Sci Pharm.* 2010;78:423-33.