

ATP depletion and oxidative damage of hepatic cells following acute exposure to malathion in rat: beneficial role of porphyrin–fullerene nanoparticles carrying magnetic magnesium

Azam Bakhtiarian¹, Mohammad Abdollahi², Seyed Mahdi Rezayat^{1,3,4}, Hamidreza Mohammadi^{5,6*}

¹Department of Pharmacology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

²Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran, Iran

³Faculty of Advanced Sciences and Technology in Medicine, Tehran University of Medical Sciences, Tehran, Iran

⁴Pharmaceutical Science Branch, Islamic Azad University, Tehran, Iran

⁵Pharmaceutical Science Research Center, Hemoglobinopathy Institute, Mazandaran University of Medical Sciences, Sari, Iran

⁶Department of Toxicology and Pharmacology, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

Received: Mar 15, 2015, Revised: May 5, 2015, Accepted: June 14, 2015

Abstract

The objective of the present study was to investigate the role of nanocarrier of magnetic isotope of $^{25}\text{Mg}^{2+}$ (PMC16) in liver toxicity, ATP content and oxidative stress due to malathion (MAL) exposure. PMC16 nanoparticles were administered in different doses (0.05, 0.1 and 0.2 LD_{50}) intravenously (iv) 40 minutes after a single MAL (0.25 LD_{50} = 207 mg/kg) intraperitoneal (ip) injection as a complement to standard therapy of pralidoxime (PAM) and atropine (AT). MgSO_4 was used as another supplement for comparison with PMC16. ATP/ADP ratio, antioxidant enzymes including catalase (CAT), glutathione peroxidase (GPx), superoxide dismutase (SOD), and oxidative stress biomarker including lipid peroxidation (LPO) were evaluated in liver tissue cells. Results indicated that after MAL administration, ADP/ATP ratio had a significant increase in liver tissues in comparison with control but this ratio was improved using various doses of PMC16. LPO was significantly decreased at all doses of PMC16 and MgSO_4 when compared with MAL-exposed group. SOD and CAT activities significantly were increased in MAL-treated group as compared to saline group. SOD was reduced by all doses of PMC16 and CAT activity was decreased in PMC16-0.1 group. These results lead us to conclude that PMC16 and MgSO_4 are so useful for protection against MAL-induced liver toxicity, ATP depletion and oxidative damages.

Keywords: Organophosphate, malathion, magnesium, nanocarrier, adenosine triphosphate, oxidative stress

Pharm Biomed Res 2015; 1(2): 10-19

DOI: 10.18869/acadpub.pbr.1.2.10

Introduction

The organophosphate compounds (OPs) are among the most widely used pesticides in the world and thus is the cause of acute and chronic toxicities in human. Mitochondria are one of the targets of OPs toxicity as recently reviewed (1). The mitochondrial

toxicity mostly appears by a reduction in cellular production of energy meaning adenosine triphosphate (ATP) that finally results in cell death. There is evidence that OP-induced hypoxia cause the induction of mitochondrial dysfunction because of the

* E-mail: hmohammadi@farabi.tums.ac.ir

severe cellular damage, followed by bioenergetics failure which leads to necrotic cell death because of ATP depletion. Also, OPs induce cholinergic receptor-independent necrotic cell death by depressing the activity of complexes I to IV in the mitochondrial respiratory chain, eliciting decrease in mitochondrial transmembrane potential, depleting intracellular ATP contents, increasing lactate dehydrogenase release and damaging cell membrane integrity (2-4). It was determined that OPs produce oxidative stress in different tissues through the formation of reactive oxygen species (ROS) and induction of oxidative stress (5-8). Hypoxia condition and oxidative stress are in favor of apoptosis and necrosis in liver tissue. Previous investigations reported that OPs induced biochemical and histopathological changes in different tissues such as liver. (9, 10) The mechanism by which OPs induce liver toxicity has not been precisely illustrated and it is difficult to identify one individual mechanism as the cause of liver toxicity related to OPs. In recent years, studies have shown that oxidative stress is one of the main molecular mechanisms of toxicity of OPs. The liver is principally sensitive to this oxidative damage due to high oxidative metabolism and a very high cytochrome P450 activity related to the xenobiotics biotransformation process and hypoxia simultaneously.

Recently, the benefit of magnesium sulfate (MgSO_4) in the management of OP poisoning was reported by our team (11). Mg is essential for ATP activity including calcium current across and within cell membranes for cardiovascular tissues. Also, the benefit of Mg-25 ($^{25}\text{Mg}^{2+}$) carrying by a nanoparticle (PMC16) against toxicity of OPs in the heart cells (12) and cardiac activity was indicated (13). Magnetic magnesium isotope ($^{25}\text{Mg}^{2+}$) carrying by a

nanoparticle called PMC16 which was previously introduced to have positive effects on production of cellular ATP (14-16).

In the present study we aimed to evaluate the situation of liver in exposure to malathion (MAL), a commonly used OP in induction of liver toxicity, and to compare the protective effects of magnesium in the ordinary form of ^{24}Mg and in the form of PMC16. For this purpose we assessed the oxidative stress biomarkers and ADP/ATP ratio in the liver tissues.

Materials and methods

Chemicals

MAL (Shimi-Keshavarz Pesticide Production Co., Tehran), pralidoxime (PAM) from (SERB, Paris), NADPH, glutathione reductase, glutathione, glutathione peroxidase, sodium azide, adenosine diphosphate (ADP) sodium salt, adenosine triphosphate (ATP) disodium salt from Sigma-Aldrich Chemie (GmbH, Munich, Germany), Tris base, methanol (HPLC-grade), malonedialdehyde (MDA), trichloroacetic acid (TCA), potassium hydroxide, diethyl ether, tetrabutylammonium hydroxide (TBAHS), MgSO_4 , n-butanol, 2-thiobarbituric acid (TBA), KH_2PO_4 (analytical grade), and H_2O_2 from Merck Chemical Co. (Tehran), and (^{25}Mg) PMC16 from Semenov institute (Russian Academy of Sciences), 2,7-dichlorodihydrofluorescein diacetate (DCFDA) from (Invitrogen), SUPELCOSIL™ LC-18-T HPLC column from SUPELCO (UK) and SOD (superoxide dismutase) kit from Randox (UK) were used in this study.

Animals and treatments

All experiments were performed with male Wistar rats weighing 200-250 g obtained

from Institute Pasteur of Tehran. The animals were housed in plastic cages in the temperature room (22 ± 2 °C), with a 12 h light/dark cycle and fed with standard rat chow (Dampars, Iran). Following several days of acclimatization, they were randomly divided into eight groups of six animals. All animal experiments were carried out in agreement with the guidelines for the care and use of laboratory animals accepted by the Japanese Pharmacological Society and acts of the Ethical Committee of Tehran University of Medical Sciences. MAL (207 mg/kg equals 0.25 LD₅₀) was dissolved in saline and administered in a constant volume of 1 ml/kg, intraperitoneally (ip). Stock solution of PMC16 was dissolved in saline (pH was adjusted at 7.86) and administered intravenous (iv) at doses of 44.8 mg/kg (0.05 LD₅₀), 89.6 mg/kg (0.1 LD₅₀), and 179.2 mg/kg (0.2 LD₅₀). Atropine (AT) was dissolved in saline and administered ip at dose of (40 mg/kg). PAM was dissolved in saline and administered ip at dose of (40 mg/kg). MgSO₄ (7H₂O) was dissolved in saline and administered iv at dose of (600 mg/kg). Control animals received only saline at an appropriate volume.

The rats were randomly divided into 8 groups as follows:

Group 1: received saline alone named Saline;

Group 2: received only MAL named MAL;

Group 3: received MAL+AT+PAM+MgSO₄ named MgSO₄;

Group 4: received MAL+AT+PAM+PMC16 (0.05 LD₅₀) named PMC16-0.05;

Group 5: received MAL+AT+PAM+PMC16 (0.1 LD₅₀) named PMC16-0.1;

Group 6: received MAL+AT+PAM+PMC16 (0.2 LD₅₀) named PMC16-0.2;

Group 7: received MAL+AT+PAM named AT+PAM;

Group 8: received MAL+PMC16 (0.05 LD₅₀) named only PMC16-0.05;

After 24 hours of the treatment, rats were anesthetized by administration of ether and underwent laparotomy. Liver tissues were dissected out and rinsed in ice-cold saline to remove the blood and immediately frozen and stored at -80 °C for various assays. Before analyzing, tissues were homogenized in 50 mM phosphate buffer solution (pH 7.4) using a tissue homogenizer at 4 °C. The homogenates were centrifuged at 3500×g for 20 min and the supernatant was used for analyses.

Measurement of lipid peroxidation (LPO)

LPO was measured by the reaction of TBA with MDA and other lipid peroxides. Samples were mixed with TCA (20%) and produced precipitate which was dispersed in H₂SO₄ (0.05 M). After addition of TBA (0.2% in sodium sulfate), the sample was heated for 30 min in a boiling water bath. Then LPO adducts were extracted by n-butanol and absorbance was measured with spectrophotometer UV 160-A (Shimadzu Japan) at 532 nm (13).

Measurement of catalase (CAT)

CAT activity was measured at 20 °C by monitoring the decomposition of 19 mM hydrogen peroxide solution prepared in potassium phosphate buffer. The disappearance of H₂O₂ was monitored at 240 nm in 1 min intervals for 5 min. Specific activity of the enzyme was expressed as micromoles per milligram protein as described previously (12)

Measurement of glutathione peroxidase (GPx)

The reaction that GPx catalyzes is: $2\text{GSH} + \text{H}_2\text{O}_2 \rightarrow \text{GS-SG} + 2\text{H}_2\text{O}$, where GSH represents reduced monomeric glutathione, and GS-SG represents glutathione disulfide. Glutathione reductase then reduces the oxidized glutathione to complete the cycle: $\text{GS-SG} + \text{NADPH} + \text{H}^+ \rightarrow 2\text{GSH} + \text{NADP}^+$. The assay mixture contained of 0.1 M phosphate buffer (pH 7.4), 1 mM GSH, 1 mM EDTA, 0.2 mM NADP (H), 1 mM sodium azide, 0.25 mM H_2O_2 and 1 mg of homogenized tissues. NADP (H) oxidation was evaluated spectrophotometrically at 340 nm.

Measurement of superoxide dismutase (SOD)

According to the kit method, xanthine and xanthine oxidase produce superoxide radicals which react with 2-(4-iodophenyl)-3-(4-nitrophenol)-5-phenyltetrazolium chloride (INT) to form a red formazan dye. The SOD activity was calculated by the degree of inhibition of this reaction in homogenized tissues. One unit of SOD causes a 50% inhibition of the rate of reduction of INT under the conditions of the assay. Data were presented as unit/mg protein (12).

Measurement of ADP and ATP

For the evaluation of the ATP/ADP ratio, tissues were sonicated in 250 μL of trichloroacetic acid (TCA) (6%) and centrifuged at 12,000g for 10 min at 4 °C. The supernatant was detached and neutralized with KOH (4 M). The cleaning of samples was done using a solid phase extraction (SPE) cartridge (C18, 22%, 100 mg/1 ml). Detection was achieved by HPLC system (Waters Chromatography Division, Milford, MA, USA) consisted of Waters 510 pump and solvent delivery system, column (SUPELCOSIL™ LC-18-

T) with guard column holder and Waters 486 ultraviolet-visible detector. The flow rate was 1 ml/min for 20 min and detection was at 254 nm. The levels of ATP and ADP were quantified after creating of standard curve, and the ADP/ATP ratio was calculated (18).

Measurement of protein

Samples were mixed with Bradford reagent dye and after 5 min, the absorbance was measured at 595 nm by the spectrophotometer using bovine serum albumin as a standard.

Statistical analysis

All data are expressed as mean \pm SEM. Statistical analysis was performed with one-way ANOVA followed by Tukey post hoc test for multiple comparisons. $P < 0.05$ was considered significant.

Results

ADP/ATP ratio

The effect of PMC16 on *ADP/ATP ratio* was investigated by assessing ADP and ATP levels in homogenated tissues of rat liver. MAL significantly increased ADP/ATP ratio in liver tissues as compared with saline group ($P < 0.05$). The ADP/ATP ratios decreased significantly in three doses of PMC16 and MgSO_4 groups as compared with MAL group (Fig. 1).

LPO

LPO level was measured by assessing TBARS level in homogenated of rat liver tissues. TBARS levels were significantly increased in the MAL-exposed group ($P < 0.001$) in comparison to saline group in liver tissues (Fig. 2). LPO was significantly decreased in only- PMC16 ($P < 0.05$), PMC16-0.05 ($P < 0.05$), PMC16-0.1 ($P < 0.05$), PMC16-0.2 ($P < 0.001$), and MgSO_4 ($P < 0.01$) in liver tissues when compared to the MAL group.




Figure 1 Effect of therapy on ADP/ATP ratio in liver tissues. Data are mean \pm SE of 6 animals in each group. Significantly different from saline group at $P < 0.05$ (a). Significantly different from MAL group at $P < 0.05$ (b).

CAT

Fig. 3 represents the effect of doses of PMC16 on CAT activity in liver rat tissues. CAT activity significantly increased ($P < 0.01$) in MAL group in comparison to Saline group. A significant decrease in the CAT activity was found in the PMC16-0.1 group ($P < 0.01$) in comparison to MAL group.

SOD

A significant increase in the SOD activity was found in Mal group as compared to saline group ($P < 0.05$) (Fig. 4). SOD activity was significantly decreased in PMC16-0.05 ($P < 0.01$), PMC16-0.1 ($P < 0.05$), PMC16-0.2 ($P < 0.001$) and even in only PMC16 ($P < 0.05$) groups in comparison to MAL group.

GPx

As shown in Fig. 5, no significant changes were found in GPx activity in different experimental groups.

Discussion

The results of this study indicate that MAL in acute exposure at a dose of 0.25 LD50 significantly increased liver ADP/ATP ratio and PMC16 in all doses used, surprisingly compensated the need of liver to ATP via increasing of ATP in hepatocytes. In addition, PMC16 reduced significantly MAL-induced LPO.

Previous studies demonstrated that OPs induce hypoxia, bradycardia, and mitochondrial dysfunction followed by depletion of cellular energy and death. This effect of MAL is mediated by inhibition of mitochondrial complexes I to IV and the respiratory chain and damaging cell membrane integrity (3, 19, 20). Nowadays, a membranotropic new magnetic magnesium ($^{25}\text{Mg}^{2+}$) carrying by fullerene nanoparticles with low toxicity has been developed. This nanoparticle, which contains obvious cationite properties, is the iron containing porphyrin monoadduct of a classical buckminster fullerene named 'Porphylleren-MC16' or briefly PMC16. In




Figure 2 Changes in lipid peroxidation as thiobarbituric acid reactive substance (TBARS) in liver tissues. Data are mean± SE of 6 animals in each group. Significantly different from Saline group at P < 0.001 (aaa).Significantly different from malathion (MAL) group at P < 0.001 (bbb), P < 0.01 (bb), P < 0.05 (b).




Figure 3 Changes in Catalase Activity in liver tissues in various groups. Data are mean ± SE of 6 animals in each group. Significantly different from saline group at P < 0.05 (a). Significantly different from malathion (MAL) group at P < 0.05 (b).




Figure 4 Changes in SOD activity in liver tissues in various groups. Data are mean \pm SE of 6 animals in each group. Significantly different from saline group at $P < 0.05$ (a). Significantly different from MAL group at $P < 0.001$ (bbb), $P < 0.01$ (bb) $P < 0.05$ (b).




Figure 5 Changes in GPx Activity in liver tissues in various groups. Data are mean \pm SE of 6 animals in each group.

acidosis condition these nanoparticles release the magnetic isotopes (^{25}Mg) and activate both oxidative phosphorylation and substrate pathways and creatine kinase (CK)-directed ADP phosphorylation inside the mitochondrial membranes in the cells (21).

According to previous studies, PMC16 is designed for a targeted delivery to compensate the ATP losses due to hypoxia-caused owing to a known ^{25}Mg -related overactivation of both oxidative and substrate ADP phosphorylation pathways of the mitochondrial . ATP synthesis by $^{25}\text{Mg}^{2+}$, the only magnetic magnesium isotope, is activated up to 2.5-fold more efficiently as compared to non-magnetic $^{24}\text{Mg}^{2+}$ and $^{26}\text{Mg}^{2+}$ isotopes. Therefore, the acidosis-induced cause release of $^{25}\text{Mg}^{2+}$ from the (^{25}Mg) PMC16 cations towards the cells and tissues suffering from any type of hypoxia in vivo administration (21, 22). Moreover, a high level of synergism manifested between the CK activity, magnesium isotopes content and ATP production rate in mitochondria and tissue acidosis degree by the PMC16 hypoxia treated cases. It should be mentioned that PMC16 does not have more affinity to hepatocytes but in this study PMC16 has increased ATP production in liver cells and prevented the disturbance in the cells by balancing their energy with a negligible amount in treated groups. This study revealed that PMC16 also has a membrane protective and LPO inhibitory effects in liver tissues that confirm our previous studies (20, 23). Demonstrated that OPs induce oxidative stress and ROS generation is one of the important of molecular mechanism of toxicity. Induced oxidative stress by OPs causes mitochondrial energy depletion (ATP depletion), induction of proteolytic enzymes and DNA fragmentation which consequences lead to necrosis or apoptosis. ROS formation can

be due to decrease rate of ATP synthesis in the mitochondria (3).

In this research although antioxidant enzymes were increased after MAL administration in MAL group as compared to saline group, PMC16 declined the activity of SOD in all used doses and CAT changed only in PMC16-0.1group. The first thing that comes to mind is the increased antioxidant power of liver reflects an activation of the compensatory mechanism through the effect of MAL on progenitor cells with its extent depending on the magnitude of the oxidative stress and therefore, on the dose of the stressor. Supporting this idea, there is evidence that administration of MAL increases the activities of CAT and SOD in erythrocytes and liver (8). No significant changes of the liver antioxidant enzymes may be due to the acute nature of the present model of the MAL poisoning or can supports this idea that PMC16 has free radical scavenger effects which this mechanism may be rather than an indirect antioxidant enzymes activation. Regarding, better efficacy of some doses of PMC16 than normal magnesium (^{24}Mg in MgSO_4) was seen in this study. It is prominent to affirm that the exceptional structure of PMC16 allows it to provide as a nanocationite with capability of releasing magnetic magnesium into hypoxic cells, in that way cause considerable enhancement of the oxidative phosphorylation pathway. Moreover, PMC16 has the porphyrin domain which provides a tissue selective interaction for porphyrin receptors located in mitochondrial membrane (24). In supporting of the present study, Mg has been found valuable in human OPs poisoning management by modulation of oxidative stress condition and reducing of the cellular ROS formation (11). Efficacy of PMC16 compared with other forms of Mg supports this idea that ^{25}Mg plays an

essential role in Mg-dependent production of ATP which consistent with the previous findings (14, 15, 23). Therefore, the PMC16 activity is dependent on delivery of ²⁵Mg isotope towards the hypoxia distressing cells and tissues.

Another hypothesis is that PMC16 molecules or its metabolites may react with MAL molecule and neutralize its electrophilic sites. Although in our previous studies we have stated that antioxidant activity was decreased, and oxidant markers were increased in malathion toxicity, the studies on the relationship between oxidant and antioxidant status and the liver damage in acute MAL poisoning are limited (6, 7, 23).

In conclusion, it is noteworthy that PMC16 is able to decrease OP-induced liver toxicity by prevention of ROS formation and elevation of ATP content in hepatic cells.

References

1. Abdollahi M, Karami-Mohajeri S. A comprehensive review on experimental and clinical findings in intermediate syndrome caused by organophosphate poisoning. *Toxicol Appl Pharm* 2012;258:309-14.
2. Karami-Mohajeri, S. Abdollahi, M. Mitochondrial dysfunction and organophosphorus compounds. *Toxicol Appl Pharm* 2013;270:39-44.
3. Bagchi D, Bagchi M, Hassoun E, Stohs S. In vitro and in vivo generation of reactive oxygen species, DNA damage and lactate dehydrogenase leakage by selected pesticides. *Toxicol* 1995;104:129-40.
4. Chan JY, Chan SH, Dai K, Cheng HL, Chou JL, Chang AY. Cholinergic receptor-independent dysfunction of mitochondrial respiratory chain enzymes, reduced mitochondrial transmembrane potential and ATP depletion underlie necrotic cell death induced by the organophosphate poison mevinphos. *Neuropharmacology* 2006;51:1109-19.
5. Banerjee B, Seth V, Bhattacharya A, Pasha S, Chakraborty A. Biochemical effects of some pesticides on lipid peroxidation and free-radical scavengers. *Toxicology lett* 1999;107:33-47.
6. Abdollahi M, Mostafalou S, Pournourmohammadi S, Shadnia S. Oxidative stress and cholinesterase inhibition in saliva and plasma of rats following subchronic exposure to malathion. *Comparative Biochemistry and Physiology Part C. Toxicol Pharm* 2004;137:29-34.
7. Abdollahi M, Ranjbar A, Shadnia S, Nikfar S, Rezaie A. Pesticides and oxidative stress: a review. *Med Sci Monitor* 2004;10:141-7
8. Akhgari M, Abdollahi M, Kebryaezadeh A, Hosseini R, Sabzevari O. Biochemical evidence for free radical-induced lipid peroxidation as a mechanism for subchronic toxicity of malathion in blood and liver of rats. *Hum Exp Toxicol* 2003;22:205-11.
9. Orchard CH, Kentish JC. Effects of changes of pH on the contractile function of cardiac muscle. *Am J Physiol* 1990;258:967-81.
10. Sutcu R, Altuntas I, Yildirim B, Karahan N, Demirin H, Delibas N. The effects of subchronic methidathion toxicity on rat liver: role of antioxidant vitamins C and E. *Cell Biol Toxicol* 2006;22:221-7.
11. Pajoumand A, Shadnia S, Rezaie A, Abdi M, Abdollahi M. Benefits of magnesium sulfate in the management of acute human poisoning by organophosphorus insecticides. *Hum Exp Toxicol* 2004;23:565-9.
12. Shafiee H, Mohammadi H, Rezayat SM, Hosseini A, Baeeri M, Hassani S, et al. Prevention of malathion-induced depletion of cardiac cells mitochondrial energy and free radical damage by a magnetic magnesium-carrying nanoparticle. *Toxicol Mech Meth* 2010;20:538-43.
13. Mohammadi H, Karimi G, Rezayat SM, Reza A, Shafiee H, Nikfar S, et al. Benefit of nanocarrier of magnetic magnesium in rat malathion-induced toxicity and cardiac failure using non-invasive monitoring of electrocardiogram and blood pressure. *Toxicol Ind Health* 2011;27:417-29.
14. Buchachenko A, Kouznetsov D, Arkhangelsky S, Orlova M, Markarian A. Spin biochemistry: Magnetic ²⁴Mg-²⁵Mg-²⁶Mg isotope effect in mitochondrial ADP phosphorylation. *Cell Biochem Biophys* 2005;43:243-51.

Acknowledgements

The authors thank Dr Dmitry A. Kuznetsov from the Department of Medicinal Nanobiotechnologies of N.I. Pirogoff Russian State Medical University for reading the paper and giving comments. The authors also thank Shokoufeh Hassani, Hoda Shafiee, Asieh Hosseini, Maryam Baeeri and Marjan Shariatpanahi for excellent technical support.

Funding

This work was supported in part by grant (#10386) from Tehran University of Medical Sciences.

Conflict of interest

The authors declare that they have no competing interests.

15. Buchachenko A, Kouznetsov D, Orlova M, Markarian A. Magnetic isotope effect of magnesium in phosphoglycerate kinase phosphorylation. *Proc Natl Acad Sci USA* 2005;102:10793-6.
16. Buchachenko A, Kuznetsov D, Arkhangel'sky S, Orlova M, Markaryan A, Berdieva A, et al. Dependence of Mitochondrial ATP Synthesis on the Nuclear Magnetic Moment of Magnesium Ions. *Dokl Biochem Biophys* 2004;396:197-9
17. Sarkhail P, Rahmanipour S, Fadyevatan S, Mohammadirad A, Dehghan G, Amin G, et al. Antidiabetic effect of *Phlomis anisodonta*: Effects on hepatic cells lipid peroxidation and antioxidant enzymes in experimental diabetes. *Pharmacol Res* 2007;56:261-6.
18. Hosseini A, Sharifzadeh M, Rezayat S. Benefit of magnesium-25 carrying porphyrin-fullerene nanoparticles in experimental diabetic neuropathy. *Int J Nanomed* 2010;5:517-23.
19. Chan J, Chan S, Dai K, Cheng H, Chou J, Chan A. Cholinergic receptor-independent dysfunction of mitochondrial respiratory chain enzymes, reduced mitochondrial transmembrane potential and ATP depletion underlie necrotic cell death induced by the organophosphate poison mevinphos. *Neuropharmacology* 2006;51:1109-19.
20. Shafiee H, Mohammadi H, Rezayat S, Hosseini A, Baeeri M, Hassani S, et al. Prevention of malathion-induced depletion of cardiac cells mitochondrial energy and free radical damage by a magnetic magnesium-carrying nanoparticle. *Toxicol Mech Meth* 2010; 20:538-43.
21. Amirshahi N, Alyautdin R, Sarkar S, Rezayat S, Orlova M, Trushkov I, et al. Porphyrin-fullerene nanoparticles for treatment of hypoxic cardiopathies. *Nanotech Russia* 2008;3:611-21.
22. Buchachenko A, Kouznetsov D, Breslavskaya NN, Orlova MA. Magnesium isotope effects in enzymatic phosphorylation. *J Phys Chem B* 2008;112: 2548-56.
23. Mohammadi HR, Karimi G, Rezayat SM, Dehpour AR, Shafiee H, Nikfar S, et al. Benefit of nanocarrier of magnetic magnesium in rat malathion-induced toxicity and cardiac failure using non-invasive monitoring of electrocardiogram and blood pressure. *Toxicol Ind Health* 2011;27:417-29.
24. Motohashi N, Takahashi A, Mifune M, Saito Y. Inhibitory effects of water-soluble cationic manganese porphyrins on peroxynitrite-induced SOS response in *Salmonella typhimurium* TA4107/pSK1002. *Mut Res Fund Mol Mech Mut* 2004;554:165-74.